

A new method for solving fully fuzzy linear Bilevel programming problems

N. Safaei*, M. Saraj

Received: 17 June 2013 ; Accepted: 4 November 2013

Abstract In this paper, a new method is proposed to find the fuzzy optimal solution of fully fuzzy linear Bilevel programming (FFLBLP) problems by representing all the parameters as triangular fuzzy numbers. In the proposed method, the given FFLBLP problem is decomposed into three crisp linear programming (CLP) problems with bounded variables constraints, the three CLP problems are solved separately and by using its optimal solutions, the fuzzy optimal solution to the given FFLBLP is obtained. The proposed method is easy to understand and to apply for finding the fuzzy optimal solution of FFLBLP occurring in real life situations.

Keywords Fully Fuzzy Linear Programming Problems, Triangular fuzzy Numbers, Bilevel Programming.

1 Introduction

Fuzzy set theory has been applied to many disciplines such as control theory and management sciences, mathematical modeling and industrial applications. The concept of fuzzy mathematical programming on general level was first proposed by Tanaka et al. [1] in the framework of the fuzzy decision of Bellman and Zadeh [2]. The first formulation of fuzzy linear programming (FLP) is proposed by Zimmermann [3]. Many researchers adopted this concept for solving fuzzy linear programming problems [1-14]. However, in all of the above mentioned works, those cases of fuzzy linear programming have been studied in which not all parts of the problem were assumed to be fuzzy, e.g., only the right hand side or the objective function coefficients were fuzzy but the variables were not fuzzy.

Allahviranloo et al. [4] solved the fuzzy integer linear programming problem by reducing it into a crisp integer linear programming problem. Allahviranloo et al. [5] proposed a new method for solving fully fuzzy linear programming problems by the use of ranking function. Nasser [6] proposed a method for solving fuzzy linear programming problems by solving the classical linear programming. Lotfi et al. [7] discussed fully fuzzy linear programming problems by representing all parameters and variables as triangular fuzzy numbers. Ebrahimnejad and Nasser [8] used the complementary slackness theorem to solve fuzzy linear programming problem with fuzzy parameters without the need of a simplex tableau. Ebrahimnejad et al. [9] proposed a new primal-dual algorithm for solving linear programming

* Corresponding Author. (✉)
E-mail: n_safaei@ymail.com (N. Safaei)

N. Safaei
M.Sc, Department of Mathematics, Faculty of Mathematical Sciences and Computer, Shahid Chamran University, Ahvaz-Iran.

M. Saraj
Associate Professor, Department of Mathematics, Faculty of Mathematical Sciences and Computer, Shahid Chamran University, Ahvaz-Iran.

problems with fuzzy variables by using duality results. Nasseri and Ebrahimnejad [10] proposed a fuzzy primal simplex algorithm for solving the flexible linear programming problem. Kumar et al. [11] proposed a new method for solving fully fuzzy linear programming problems by introducing fuzzy slack and surplus variables. Kumar et al. [12, 13] proposed a new method for solving fully fuzzy linear programming problems with inequality constraints. Kumar et al. [14] proposed a new method for finding the fuzzy optimal solution of fully fuzzy linear programming problems with equality constraints.

This paper is organized as follows: In Section 2 some basic definitions and arithmetics between two triangular fuzzy numbers are reviewed. In Section 3 formulation of FLBLP problems and a new method is proposed for solving FLBLP problems we solve an illustrative numerical example in Section 4.

2 Preliminaries

In this section, some necessary backgrounds and notions of fuzzy set theory are reviewed.

2.1 Basic definitions

Definition 2.1 [15]. The characteristic function μ_A of a crisp set $A \subseteq X$ assigns a value either 0 or 1 to each member in X . This function can be generalized to a function $\mu_{\tilde{A}}$ such that the value assigned to the element of the universal set X fall within a specified range i.e. $\mu_{\tilde{A}} : X \rightarrow [0,1]$. The assigned value indicate the membership grade of the element in the set A .

The function $\mu_{\tilde{A}}$ is called the membership function and the set $\tilde{A} = \{(x, \mu_{\tilde{A}}(x)); x \in X\}$ defined by $\mu_{\tilde{A}}(x)$ for each $x \in X$ is called a fuzzy set.

Definition 2.2 [15]. A fuzzy number $\tilde{A} = (a, b, c)$ is said to be a triangular fuzzy number if its membership function is given by

$$\mu_{\tilde{A}}(x) = \begin{cases} \frac{x-a}{b-a} & a \leq x \leq b \\ 1 & x = b \\ \frac{x-c}{b-c} & b \leq x \leq c \end{cases}$$

Definition 2.3 [15]. A triangular fuzzy number (a, b, c) is said to be non-negative fuzzy number iff $a \geq 0$.

Definition 2.4 [15]. A fuzzy number \tilde{A} is said to be non-negative fuzzy number if and only if $\mu_{\tilde{A}}(x) = 0$, for all $x < 0$.

Definition 2.5 [15]. Two triangular fuzzy numbers $\tilde{A} = (a, b, c)$ and $\tilde{B} = (e, f, g)$ are said to be equal if and only if $a = e, b = f, c = g$.

2.2 Arithmetic operations

In this subsection, arithmetic operations between two triangular fuzzy numbers, defined on universal set of real numbers, are reviewed [4].

Let $\tilde{A} = (a, b, c)$ and $\tilde{B} = (e, f, g)$ be two triangular fuzzy numbers then

$$(i) \quad \tilde{A} \oplus \tilde{B} = (a, b, c) \oplus (e, f, g) = (a+e, b+f, c+g)$$

$$(ii) \quad -\tilde{A} = -(a, b, c) = (-c, -b, -a)$$

(iii) Let $\tilde{A} = (a, b, c)$ be any triangular fuzzy number and Let $\tilde{B} = (x, y, z)$ be a non-negative triangular fuzzy number then

$$\tilde{A} \otimes \tilde{B} = \begin{cases} (ax, by, cz) & a \geq 0 \\ (az, by, cz) & a < 0, c \geq 0 \\ (az, by, cx) & c < 0 \end{cases}$$

3 Fully fuzzy linear bilevel programming problem

Consider a FFLBLPP of maximization-type objective functions at each level. Suppose that DM_i denotes the DM at the i -th level ($i = 1, 2$) who controls the decision vector

$$\underset{\tilde{x}_1}{\text{Max}} \quad \tilde{Z}_1 \approx \tilde{C}_{11}^T \tilde{x}_1 \oplus \tilde{C}_{12}^T \tilde{x}_2 \oplus \tilde{\alpha}_1$$

where \tilde{x}_2 solve

$$\underset{\tilde{x}_1, \tilde{x}_2}{\text{Max}} \quad \tilde{Z}_2 \approx \tilde{C}_{21}^T \tilde{x}_1 \oplus \tilde{C}_{22}^T \tilde{x}_2 \oplus \tilde{\alpha}_2$$

s.t.

$$\tilde{A} \tilde{x}_1 \oplus \tilde{B} \tilde{x}_2 \{ \leq, \approx, \geq \} \tilde{t}$$

$$\tilde{x}_1, \tilde{x}_2 \geq 0$$

Where $\tilde{\alpha}_1$ and $\tilde{\alpha}_2$ are fuzzy number, $\tilde{A} = (\tilde{a}_{ij})_{m \times n_1}, \tilde{B} = (\tilde{b}_{ij})_{m \times n_2}, \tilde{t} = (\tilde{t}_i)_{m \times 1}$ and $\tilde{C}_{11} = (C_{11j})_{1 \times n_1}, \tilde{C}_{12} = (C_{21j})_{1 \times n_2}, \tilde{C}_{21} = (C_{21j})_{1 \times n_1}, \tilde{C}_{22} = (C_{22j})_{1 \times n_2}, n_1 + n_2 = n, \tilde{a}_{ij}, \tilde{b}_{ij}, \tilde{x}_1, \tilde{x}_2, \tilde{t}_i \in F(\mathbb{R})$ for all $1 \leq i \leq m, 1 \leq j \leq n$.

Let the parameters $\tilde{C}_{11}, \tilde{C}_{12}, \tilde{C}_{21}, \tilde{C}_{22}, \tilde{a}_{ij}, \tilde{b}_{ij}, \tilde{x}_1, \tilde{x}_2$ and \tilde{t}_i be the triangular fuzzy number $(p_{1j}, q_{1j}, r_{1j}), (p_{2j}, q_{2j}, r_{2j}), (p_{3j}, q_{3j}, r_{3j}), (p_{4j}, q_{4j}, r_{4j}), (a_{ij}^{-1}, a_{ij}^{-2}, a_{ij}^{-3}), (b_{ij}^{-1}, b_{ij}^{-2}, b_{ij}^{-3}), (x_{1j}, y_{1j}, t_{1j}), (x_{2j}, y_{2j}, t_{2j})$ and (m_i, n_i, p_i) respectively. Then, the problem can be written as follows:

$$\underset{x_{2j}, y_{2j}, t_{2j}}{\text{Max}} (Z_{1L}, Z_{1M}, Z_{1U}) = \sum_{j=1}^{n_1} (p_{1j}, q_{1j}, r_{1j}) \oplus (x_{1j}, y_{1j}, t_{1j}) \oplus \sum_{j=1}^{n_2} (p_{2j}, q_{2j}, r_{2j}) \otimes (x_{2j}, y_{2j}, t_{2j})$$

where x_{2j}, y_{2j}, t_{2j} solves

$$\text{Max } (Z_{2L}, Z_{2M}, Z_{2U}) = \sum_{j=1}^{n_1} (p_{3j}, q_{3j}, r_{3j}) \oplus (x_{1j}, y_{1j}, t_{1j}) \oplus \sum_{j=1}^{n_2} (p_{4j}, q_{4j}, r_{4j}) \otimes (x_{2j}, y_{2j}, t_{2j})$$

s.t.

$$\sum_{j=1}^{n_1} (a_{ij}^{-1}, a_{ij}^{-2}, a_{ij}^{-3}) \oplus (x_{1j}, y_{1j}, t_{1j}) \oplus \sum_{j=1}^{n_2} (b_{ij}^{-1}, b_{ij}^{-2}, b_{ij}^{-3}) \otimes (x_{2j}, y_{2j}, t_{2j}) \{ \leq, =, \geq \} (m_i, n_i, p_i)$$

for all $i = 1, \dots, m$.

and all decision variables are non-negative.

Now, using the arithmetic operations and partial ordering relations, we decompose the given FBLPP as follows:

$$\underset{x_{2j}}{\text{Max}} Z_{1L} = \text{lower value of} \left[\sum_{j=1}^{n_1} (p_{1j}, q_{1j}, r_{1j}) \oplus (x_{1j}, y_{1j}, t_{1j}) \oplus \sum_{j=1}^{n_2} (p_{2j}, q_{2j}, r_{2j}) \otimes (x_{2j}, y_{2j}, t_{2j}) \right]$$

$$\underset{y_{2j}}{\text{Max}} Z_{1M} = \text{middle value of} \left[\sum_{j=1}^{n_1} (p_{1j}, q_{1j}, r_{1j}) \oplus (x_{1j}, y_{1j}, t_{1j}) \oplus \sum_{j=1}^{n_2} (p_{2j}, q_{2j}, r_{2j}) \otimes (x_{2j}, y_{2j}, t_{2j}) \right]$$

$$\underset{t_{2j}}{\text{Max}} Z_{1U} = \text{upper value of} \left[\sum_{j=1}^{n_1} (p_{1j}, q_{1j}, r_{1j}) \oplus (x_{1j}, y_{1j}, t_{1j}) \oplus \sum_{j=1}^{n_2} (p_{2j}, q_{2j}, r_{2j}) \otimes (x_{2j}, y_{2j}, t_{2j}) \right]$$

where x_{2j}, y_{2j}, t_{2j} solves

$$\text{Max } Z_{2L} = \text{lower value of} \left[\sum_{j=1}^{n_1} (p_{3j}, q_{3j}, r_{3j}) \oplus (x_{1j}, y_{1j}, t_{1j}) \oplus \sum_{j=1}^{n_2} (p_{4j}, q_{4j}, r_{4j}) \otimes (x_{2j}, y_{2j}, t_{2j}) \right]$$

$$\text{Max } Z_{2M} = \text{middle value of} \left[\sum_{j=1}^{n_1} (p_{3j}, q_{3j}, r_{3j}) \oplus (x_{1j}, y_{1j}, t_{1j}) \oplus \sum_{j=1}^{n_2} (p_{4j}, q_{4j}, r_{4j}) \otimes (x_{2j}, y_{2j}, t_{2j}) \right]$$

$$\text{Max } Z_{2U} = \text{upper value of} \left[\sum_{j=1}^{n_1} (p_{3j}, q_{3j}, r_{3j}) \oplus (x_{1j}, y_{1j}, t_{1j}) \oplus \sum_{j=1}^{n_2} (p_{4j}, q_{4j}, r_{4j}) \otimes (x_{2j}, y_{2j}, t_{2j}) \right]$$

s.t.

$$\text{lower value of} \left[\sum_{j=1}^{n_1} (a_{ij}^{-1}, a_{ij}^{-2}, a_{ij}^{-3}) \oplus (x_{1j}, y_{1j}, t_{1j}) \oplus \sum_{j=1}^{n_2} (b_{ij}^{-1}, b_{ij}^{-2}, b_{ij}^{-3}) \otimes (x_{2j}, y_{2j}, t_{2j}) \{ \leq, =, \geq \} m_i \right]$$

$$\text{middle value of} \left[\sum_{j=1}^{n_1} (a_{ij}^{-1}, a_{ij}^{-2}, a_{ij}^{-3}) \oplus (x_{1j}, y_{1j}, t_{1j}) \oplus \sum_{j=1}^{n_2} (b_{ij}^{-1}, b_{ij}^{-2}, b_{ij}^{-3}) \otimes (x_{2j}, y_{2j}, t_{2j}) \{ \leq, =, \geq \} n_i \right]$$

$$\text{upper value of} \left[\sum_{j=1}^{n_1} (a_{ij}^{-1}, a_{ij}^{-2}, a_{ij}^{-3}) \oplus (x_{1j}, y_{1j}, t_{1j}) \oplus \sum_{j=1}^{n_2} (b_{ij}^{-1}, b_{ij}^{-2}, b_{ij}^{-3}) \otimes (x_{2j}, y_{2j}, t_{2j}) \{ \leq, =, \geq \} p_i \right]$$

for all $i = 1, \dots, m$.

and all decision variables are non-negative.

From the above decomposition problem, we construct the following CLP problems namely, middle level bilevel problem (MLBLP), upper level bilevel problem (ULBLP) and lower level bilevel problem (LLBLP) as follows:

(MLBLP)

$$\underset{y_{2j}}{\text{Max}} Z_{1M} = \text{middle value of} \left[\sum_{j=1}^{n_1} (p_{1j}, q_{1j}, r_{1j}) \oplus (x_{1j}, y_{1j}, t_{1j}) \oplus \sum_{j=1}^{n_2} (p_{2j}, q_{2j}, r_{2j}) \otimes (x_{2j}, y_{2j}, t_{2j}) \right]$$

where y_{2j} solve

$$\underset{x_{2j}}{\text{Max}} Z_{2M} = \text{middle value of} \left[\sum_{j=1}^{n_1} (p_{3j}, q_{3j}, r_{3j}) \oplus (x_{1j}, y_{1j}, t_{1j}) \oplus \sum_{j=1}^{n_2} (p_{4j}, q_{4j}, r_{4j}) \otimes (x_{2j}, y_{2j}, t_{2j}) \right]$$

s.t.

Constraints in the decomposition problem in which at least one decision variable of the MLBLP occurs and all decision variables are non-negative.

(LLBLP)

$$\underset{x_{2j}}{\text{Max}} Z_{1L} = \text{lower value of} \left[\sum_{j=1}^{n_1} (p_{1j}, q_{1j}, r_{1j}) \oplus (x_{1j}, y_{1j}, t_{1j}) \oplus \sum_{j=1}^{n_2} (p_{2j}, q_{2j}, r_{2j}) \otimes (x_{2j}, y_{2j}, t_{2j}) \right]$$

where x_{2j} solve

$$\underset{x_{2j}}{\text{Max}} Z_{2L} = \text{lower value of} \left[\sum_{j=1}^{n_1} (p_{3j}, q_{3j}, r_{3j}) \oplus (x_{1j}, y_{1j}, t_{1j}) \oplus \sum_{j=1}^{n_2} (p_{4j}, q_{4j}, r_{4j}) \otimes (x_{2j}, y_{2j}, t_{2j}) \right]$$

s.t.

$$x_{1j} \leq y_{1j}^*, x_{2j} \leq y_{2j}^*$$

Constraints in the decomposition constraints in which at least one decision variable of the LLBLP occurs which are not used in MLBLP and ULBLP; all variables in the constraints and objective function in LLBLP must satisfy the bounded constraints; replacing all values of the decision variables which are obtained in the MLBLP and all decision variables are non-negative. Where y_{1j}^*, y_{2j}^* is the optimal value of MLBLP.

(ULBLP)

$$\underset{t_{2j}}{\text{Max}} Z_{1U} = \text{upper value of} \left[\sum_{j=1}^{n_1} (p_{1j}, q_{1j}, r_{1j}) \oplus (x_{1j}, y_{1j}, t_{1j}) \oplus \sum_{j=1}^{n_2} (p_{2j}, q_{2j}, r_{2j}) \otimes (x_{2j}, y_{2j}, t_{2j}) \right]$$

where t_{2j} solve

$$\underset{t_{2j}}{\text{Max}} Z_{2U} = \text{upper value of} \left[\sum_{j=1}^{n_1} (p_{3j}, q_{3j}, r_{3j}) \oplus (x_{1j}, y_{1j}, t_{1j}) \oplus \sum_{j=1}^{n_2} (p_{4j}, q_{4j}, r_{4j}) \otimes (x_{2j}, y_{2j}, t_{2j}) \right]$$

s.t.

$$t_{1j} \geq y_{1j}^*, t_{2j} \geq y_{2j}^*$$

Constraints in the decomposition problem in which atleast one decision variable of the ULBLP occurs and are not used in MLBLP; all variables in the constraints and objective function in ULBLP must satisfy the bounded constraints ; replacing all values of the decision variables which are obtained in MLBLP

and LLBLP and all decision variables are non-negative. Where y_{1j}^*, y_{2j}^* is the optimal value of MLBLP.

4 Numerical example

Example : Consider the following fully fuzzy linear Bilevel programming problem:

$$\underset{\tilde{x}_1}{\text{Max}} \quad \tilde{Z}_1 \approx (3,5,7) \otimes \tilde{x}_1 \oplus (2,4,8) \otimes \tilde{x}_2$$

where \tilde{x}_2 solve

$$\underset{\text{s.t.}}{\text{Max}} \quad \tilde{Z}_2 \approx (3,5,10) \otimes \tilde{x}_1 \oplus (1,7,8) \otimes \tilde{x}_2$$

$\quad (4,5,9) \otimes \tilde{x}_1 \oplus (2,7,8) \otimes \tilde{x}_2 \leq (4,10,20)$

$$\quad (0,3,7) \otimes \tilde{x}_1 \oplus (1,2,10) \otimes \tilde{x}_2 \leq (2,5,18)$$

$$\quad \tilde{x}_1, \tilde{x}_2 \geq 0$$

Now, the decomposition problem of the given FLPP is given below:

$$\underset{\text{Max}}{\text{Max}} \quad Z_{1L} = 3x_1 + 2x_2$$

$$\underset{\text{Max}}{\text{Max}} \quad Z_{1M} = 5y_1 + 4y_2$$

$$\underset{\text{Max}}{\text{Max}} \quad Z_{1U} = 7t_1 + 8t_2$$

where x_2, y_2 and t_2 solves

$$\underset{\text{Max}}{\text{Max}} \quad Z_{2L} = 3x_1 + x_2$$

$$\underset{\text{Max}}{\text{Max}} \quad Z_{2M} = 5y_1 + 7y_2$$

$$\underset{\text{Max}}{\text{Max}} \quad Z_{2U} = 10t_1 + 8t_2$$

s.t.

$$4x_1 + 2x_2 \leq 4, 5y_1 + 7y_2 \leq 10, 9t_1 + 8t_2 \leq 20$$

$$x_2 \leq 2, 3y_1 + 2y_2 \leq 5, 7t_1 + 10t_2 \leq 18$$

$$x_i, y_i, t_i \geq 0, \quad i = 1, 2.$$

Now, the Middle Level problem is given below:

(MLBLP)

$$\underset{\text{Max}}{\text{Max}} \quad Z_{1M} = 5y_1 + 4y_2$$

where y_2 solve

$$\underset{\text{Max}}{\text{Max}} \quad Z_{2M} = 5y_1 + 7y_2$$

s.t.

$$5y_1 + 7y_2 \leq 10, 3y_1 + 2y_2 \leq 5$$

$$y_1, y_2 \geq 0$$

Now, solving the problem(MLBLP)using simplex method, we obtain the optimal solution $y_1 = 1.36$, $y_2 = 0.45$ and $Z_{1M} = 8.6$.

Now, the lower Level problem is given below:

(LBLLP)

$$\text{Max } Z_{1L} = 3x_1 + 2x_2$$

where x_2 solve

$$\text{Max } Z_{2L} = 3x_1 + x_2$$

s.t.

$$4x_1 + 2x_2 \leq 4, x_2 \leq 2$$

$$x_1 \leq 1.36, x_2 \leq 0.45$$

$$x_1, x_2 \geq 0$$

Now, solving the problem LLP with $y_1 = 1.36$, $y_2 = 0.45$ by simplex method, we obtain the optimal solution $x_1 = 0.39$, $x_2 = 0.45$ and $Z_{1L} = 2.07$.

Now, the Upper Level problem is given below:

(ULBLP)

$$\text{Max } Z_{1U} = 7t_1 + 8t_2$$

where t_2 solves

$$\text{Max } Z_{2U} = 10t_1 + 8t_2$$

s.t.

$$9t_1 + 8t_2 \leq 20, 7t_1 + 10t_2 \leq 18$$

$$t_1 \geq 1.36, t_2 \geq 0.45$$

$$t_1, t_2 \geq 0$$

Now, solving the problem LLPwith $y_1 = 1.36$, $y_2 = 0.45$ by simplex method, we obtain the optimal solution $t_1 = 1.65$, $t_2 = 0.65$ and $Z_{1U} = 16.75$.

Therefore, an optimal fuzzy solution to the given fully fuzzy linear Bilevel programming problem is

$$\tilde{x}_1 = (x_1, y_1, t_1) = (0.39, 1.36, 1.65), \tilde{x}_2 = (x_2, y_2, t_2) = (0.45, 0.45, 0.45), \tilde{Z}_1 = (Z_{1L},$$

$$Z_{1M}, Z_{1U}) = (2.07, 8.6, 16.75) \text{ and } \tilde{Z}_2 = (Z_{2L}, Z_{2M}, Z_{2U}) = (1.62, 9.95, 21.7).$$

5 Conclusions

In this paper, fuzzy linear Bilevel programming problems with fuzzy variables and fuzzy constraints are discussed. The significant of this paper is solving fuzzy linear Bilevel programming problem when the variables triangular fuzzy numbers without using ranking method. Thus the method is very useful in the real world problems where the product is uncertain.

References

1. Tanaka, H., Okuda, T., Asai, K., (1974). On fuzzy mathematical programming. *Journal of Cybernetics*, 3, 37-46.
2. Bellman, R. E., Zadeh, L. A., (1970). Decision making in a fuzzy environment. *Management Science*, 17, 141-164.
3. Zimmermann, H. J., (1978). Fuzzy programming and linear programming with several objective functions. *Fuzzy Sets and Systems*, 1, 45-55.
4. Allahviranloo, T., Shamsolkotabi, K. H., Kiani, N. A., Alizadeh, L., (2007). Fuzzy integer linear programming problems. *International Journal of Contemporary Mathematical Sciences*, 2, 167-181.
5. Allahviranloo, T., Lotfi, F. H., Kiasary, M. K., Kiani, N. A., Alizadeh, L., (2008). Solving fully fuzzy linear programming problem by the ranking function. *Applied Mathematical Sciences*, 2, 19-32.
6. Nasseri, S. H., (2008). A new method for solving fuzzy linear programming by solving linear programming. *Applied Mathematical Sciences*, 2, 2473-2480.
7. Lotfi, F. H., Allahviranloo, T., Jondabeha, M. A., Alizadeh, L., (2009). Solving a fully fuzzy linear programming using lexicography method and fuzzy approximate solution, *Applied Mathematical Modelling*, 33, 3151-3156.
8. Ebrahimnejad, A., Nasseri, S. H., (2009). Using complementary slackness property to solve linear programming with fuzzy parameters, *Fuzzy Information and Engineering*, 1, 233-245.
9. Ebrahimnejad, A., Nasseri, S. H., Lotfi, F. H., Soltanifar, M., (2010). A primal-dual method for linear programming problems with fuzzy variables. *European Journal of Industrial Engineering*, 4, 189-209.
10. Nasseri, S. H., Ebrahimnejad, A., (2010). A fuzzy primal simplex algorithm and its applications for solving flexible linear programming problems. *European Journal of Industrial Engineering*, 4, 372-389.
11. Kumar, A., Kaur, J., Singh, P., (2010). Fuzzy linear programming problems with fuzzy Parameters. *Journal of Advanced Research in Scientific Computing*, 2, 1-12.
12. Kumar, A., Kaur, J., Singh, P., (2010). Solving fully fuzzy linear programming problems with inequality constraints. *International Journal of Physical and Mathematical Sciences*, 1, 6-18.
13. Kumar, A., Kaur, J., Singh, P., (2010). Fuzzy Optimal Solution of Fully Fuzzy Linear Programming Problems with Inequality Constraints. *International Journal of Applied Mathematics and Computer Sciences*, 6, 37-41.
14. Kumar, A., Kaur, J., Singh, P., (2011). A new method for solving fully fuzzy linear programming problems. *Applied Mathematical Modelling*, 35, 817-823.
15. Dubois, D., Prade, H., (1980). *Fuzzy Sets and Systems: Theory and Applications*, Academic Press, New York.