

## Optimization of solution Kadomtsev-Petviashvili equation by using homotopy methods

T. Allahviranloo\*, Sh. S. Behzadi

**Received:** 8 June 2013 ; **Accepted:** 10 December 2013

**Abstract** In this paper, the Kadomtsev-Petviashvili equation is solved by using the Adomian's decomposition method, modified Adomian's decomposition method, variational iteration method, modified variational iteration method, homotopy perturbation method, modified homotopy perturbation method and homotopy analysis method. The existence and uniqueness of the solution and convergence of the proposed methods are proved in details. A numerical example is studied to demonstrate the accuracy of the presented methods.

**Keywords** Kadomtsev-Petviashvili equation, Adomian decomposition method (ADM), Modified Adomian decomposition method (MADM), Variational iteration method (VIM), Modified variational iteration method (MVIM), Homotopy perturbation method (HPM), Modified homotopy perturbation method (MHPM), Homotopy analysis method (HAM).

### 1 Introduction

In 1970, Kadomtsev and Petviashvili [1] generalized the KDV equation to two space variables and formulated the well-known Kadomtsev-Petviashvili equation to provide an explanation of the general weakly dispersive waves [2-10]. In this work, we develop the ADM, MADM, VIM, MVIM, HPM, MHPM and HAM to solve this equation as follows:

$$u_t + \mu(x, t)u_x + \frac{1}{2}\sigma^2(x, t)u_{xx} - v(x, t)u + s(x, t) = 0. \quad (1)$$

With the initial condition:

$$u(x, 0) = g(x). \quad (2)$$

Where  $g(x)$ ,  $\mu(x, t)$ ,  $\sigma(x, t)$ ,  $v(x, t)$  and  $s(x, t)$  are known functions.

The paper is organized as follows. In section 2, the mentioned iterative methods are introduced for solving Eq.(1). In section 3 we prove the existence, uniqueness of the solution and convergence of the proposed methods. Finally, the numerical example is shown in section 4.

---

\* **Corresponding Author.** (✉)

**E-mail:** [tofigh@allahviranloo.com](mailto:tofigh@allahviranloo.com) (T. Allahviranloo)

**T. Allahviranloo**

Professor, Department of Mathematics, Science and Research Branch, Islamic Azad University, Tehran, Iran.

**Sh. S. Behzadi**

Assistant professor, Department of Mathematics, Islamic Azad University, Qazvin Branch, Qazvin, Iran.

In order to obtain an approximate solution of Eq.(1), let us integrate one time Eq.(1) with respect to  $t$  using the initial condition we obtain,

$$u(x, t) = G(x, t) - \int_0^t F_1(u(x, \tau)) d\tau - \int_0^t F_2(u(x, \tau)) d\tau + \int_0^t F_3(u(x, \tau)) d\tau, \quad (3)$$

where,

$$\begin{aligned} G(x, t) &= g(x) - \int_0^t s(x, \tau) d\tau, \\ F_1(u(x, t)) &= \mu(x, t) u_x(x, t), \\ F_2(u(x, t)) &= \frac{1}{2} \sigma^2(x, t) u_{xx}(x, t), \\ F_3(u(x, t)) &= v(x, t) u(x, t). \end{aligned}$$

In Eq.(3), we assume  $G(x, t)$  is bounded for all  $x, t$  in  $J = [0, T](T \in \mathbb{R})$ .

The terms  $F_1(u(x, t))$ ,  $F_2(u(x, t))$  and  $F_3(u(x, t))$  are Lipschitz continuous with

$$|F_1(u) - F_1(u^*)| \leq L_1 |u - u^*|, \quad |F_2(u) - F_2(u^*)| \leq L_2 |u - u^*| \quad \text{and} \quad |F_3(u) - F_3(u^*)| \leq L_3 |u - u^*|.$$

## 2 The iterative methods

### 2.1 Description of the MADM and ADM

The Adomian decomposition method is applied to the following general nonlinear equation

$$Lu + Ru + Nu = f, \quad (4)$$

where  $u(x, t)$  is the unknown function,  $L$  is the highest order derivative operator which is assumed to be easily invertible,  $R$  is a linear differential operator of order less than  $L$ ,  $Nu$  represents the nonlinear terms, and  $f$  is the source term. Applying the inverse operator  $L^{-1}$  to both sides of Eq.(4), and using the given conditions we obtain

$$u(x, t) = z(x) - L^{-1}(Ru) - L^{-1}(Nu), \quad (5)$$

where the function  $z(x)$  represents the terms arising from integrating the source term  $f$ . The nonlinear operator  $Nu = G_1(u)$  is decomposed as

$$G_1(u) = \sum_{n=0}^{\infty} A_n, \quad (6)$$

where  $A_n, n \geq 0$  are the Adomian polynomials determined formally as follows:

$$A_n = \frac{1}{n!} \left[ \frac{d^n}{d\lambda^n} \left[ N \left( \sum_{i=0}^{\infty} \lambda^i u_i \right) \right] \right]_{\lambda=0}. \quad (7)$$

The first Adomian polynomials (introduced in [11,12,13]) are:

$$\begin{aligned}
A_0 &= G_1(u_0), \\
A_1 &= u_1 G_1'(u_0), \\
A_2 &= u_2 G_1'(u_0) + \frac{1}{2!} u_1^2 G_1''(u_0), \\
A_3 &= u_3 G_1'(u_0) + u_1 u_2 G_1''(u_0) + \frac{1}{3!} u_1^3 G_1'''(u_0), \dots
\end{aligned} \tag{8}$$

### 2.1.1 Adomian decomposition method

The standard decomposition technique represents the solution of  $u(x, t)$  in (4) as the following series,

$$u(x, t) = \sum_{i=0}^{\infty} u_i(x, t), \tag{9}$$

where, the components  $u_0(x, t), u_1(x, t), \dots$  which can be determined recursively

$$\begin{aligned}
u_0(x, t) &= G(x, t), \\
u_1(x, t) &= -\int_0^t A_0(x, t) dt - \int_0^t B_0(x, t) dt + \int_0^t L_0(x, t) dt, \\
&\vdots \\
u_{n+1}(x, t) &= -\int_0^t A_n(x, t) dt - \int_0^t B_n(x, t) dt + \int_0^t L_n(x, t) dt, \quad n \geq 0.
\end{aligned} \tag{10}$$

Substituting (8) into (10) leads to the determination of the components of  $u(x, t)$ .

### 2.1.2 The modified Adomian decomposition method

The modified decomposition method was introduced by Wazwaz [14]. The modified forms was established on the assumption that the function  $G(x, t)$  can be divided into two parts, namely  $G_1(x, t)$  and  $G_2(x, t)$ . Under this assumption we set

$$G(x, t) = G_1(x, t) + G_2(x, t). \tag{11}$$

Accordingly, a slight variation was proposed only on the components  $u_0$  and  $u_1$ . The suggestion was that only the part  $G_1$  be assigned to the zeroth component  $u_0$ , whereas the remaining part  $G_2$  be combined with the other terms given in (11) to define  $u_1$ . Consequently, the modified recursive relation

$$\begin{aligned}
u_0(x, t) &= G_1(x, t), \\
u_1(x, t) &= G_2(x, t) - L^{-1}(Ru_0) - L^{-1}(A_0), \\
&\vdots \\
u_{n+1}(x, t) &= -L^{-1}(Ru_n) - L^{-1}(A_n), \quad n \geq 1,
\end{aligned} \tag{12}$$

was developed.

To obtain the approximation solution of Eq.(1), according to the MADM, we can write the iterative formula (12) as follows:

$$\begin{aligned} u_0(x, t) &= G_1(x, t), \\ u_1(x, t) &= G_2(x, t) - \int_0^t A_0(x, t) dt - \int_0^t B_0(x, t) dt + \int_0^t L_0(x, t) dt, \\ &\vdots \\ u_{n+1}(x, t) &= -\int_0^t A_n(x, t) dt - \int_0^t B_n(x, t) dt + r \int_0^t L_n(x, t) dt, \quad n \geq 1. \end{aligned} \quad (13)$$

The operators  $F_i(u(x, t))$  ( $i=1,2,3$ ) are usually represented by the infinite series of the Adomian polynomials as follows:

$$\begin{aligned} F_1(u) &= \sum_{i=0}^{\infty} A_i, \\ F_2(u) &= \sum_{i=0}^{\infty} B_i, \\ F_3(u) &= \sum_{i=0}^{\infty} L_i. \end{aligned}$$

where  $A_i$ ,  $B_i$  and  $L_i$  are the Adomian polynomials.

Also, we can use the following formula for the Adomian polynomials [15]:

$$\begin{aligned} A_n &= F_1(s_n) - \sum_{i=0}^{n-1} A_i, \\ B_n &= F_2(s_n) - \sum_{i=0}^{n-1} B_i, \\ L_n &= F_3(s_n) - \sum_{i=0}^{n-1} L_i. \end{aligned} \quad (14)$$

Where  $s_n = \sum_{i=0}^n u_i(x, t)$  is the partial sum.

## 2.2 Description of the VIM and MVIM

In the VIM [ 16-23], it has been considered the following nonlinear differential equation:

$$Lu + Nu = G, \quad (15)$$

where  $L$  is a linear operator,  $N$  is a nonlinear operator and  $G$  is a known analytical function. In this case, the functions  $u_n$  may be determined recursively by

$$u_{n+1}(x, t) = u_n(x, t) + \int_0^t \lambda(x, \tau) \{L(u_n(x, \tau)) + N(u_n(x, \tau)) - G(x, \tau)\} d\tau, \quad n \geq 0, \quad (16)$$

where  $\lambda$  is a general Lagrange multiplier which can be computed using the variational theory. Here the function  $u_n(x, \tau)$  is a restricted variations which means  $\delta u_n = 0$ . Therefore, we first determine the Lagrange multiplier  $\lambda$  that will be identified optimally via integration by parts. The successive approximation  $u_n(x, t)$ ,  $n \geq 0$  of the solution  $u(x, t)$  will be readily obtained upon using the obtained Lagrange multiplier and by using any selective function  $u_0$ . The zeroth approximation  $u_0$  may be selected any function that just satisfies at least the initial and boundary conditions. With  $\lambda$  determined, then several approximation  $u_n(x, t)$ ,  $n \geq 0$  follow immediately. Consequently, the exact solution may be obtained by using

$$u(x, t) = \lim_{n \rightarrow \infty} u_n(x, t). \quad (17)$$

The VIM has been shown to solve effectively, easily and accurately a large class of nonlinear problems with approximations converge rapidly to accurate solutions.

To obtain the approximation solution of Eq.(1), according to the VIM, we can write iteration formula (16) as follows:

$$u_{n+1}(x, t) = u_n(x, t) + L_t^{-1}(\lambda[u_n(x, t) - G(x, t) + \int_0^t F_1(u_n(x, t)) dt + \int_0^t F_2(u_n(x, t)) dt - \int_0^t F_2(u_n(x, t)) dt]), n \geq 0. \quad (18)$$

Where,

$$L_t^{-1}(\cdot) = \int_0^t (\cdot) d\tau.$$

To find the optimal  $\lambda$ , we proceed as

$$\delta u_{n+1}(x, t) = \delta u_n(x, t) + \delta L_t^{-1}(\lambda[u_n(x, t) - G(x, t) + \int_0^t F_1(u_n(x, t)) dt + \int_0^t F_2(u_n(x, t)) dt - \int_0^t F_3(u_n(x, t)) dt]). \quad (19)$$

From Eq.(19), the stationary conditions can be obtained as follows:

$$\lambda' = 0 \quad \text{and} \quad 1 + \lambda = 0.$$

Therefore, the Lagrange multipliers can be identified as  $\lambda = -1$  and by substituting in (18), the following iteration formula is obtained.

$$u_0(x, t) = G(x, t), \\ u_{n+1}(x, t) = u_n(x, t) - L_t^{-1}(u_n(x, t) - G(x, t) + \int_0^t F_1(u_n(x, t)) dt + \int_0^t F_2(u_n(x, t)) dt - \int_0^t F_3(u_n(x, t)) dt), n \geq 0. \quad (20)$$

To obtain the approximation solution of Eq.(1), based on the MVIM [24,25], we can write the following iteration formula:

$$\begin{aligned}
u_0(x, t) &= G(x, t), \\
u_{n+1}(x, t) &= u_n(x, t) - L_t^{-1} \left( \int_0^t F_1(u_n(x, t) - u_{n-1}(x, t)) dt \right. \\
&\quad \left. + \int_0^t F_2(u_n(x, t) - u_{n-1}(x, t)) dt - \int_0^t F_3(u_n(x, t) - u_{n-1}(x, t)) dt \right), n \geq 0.
\end{aligned} \tag{21}$$

Relations (20) and (21) will enable us to determine the components  $u_n(x, t)$  recursively for  $n \geq 0$ .

### 2.3 Description of the HAM

Consider  $N[u] = 0$ , where  $N$  is a nonlinear operator,  $u(x, t)$  is an unknown function and  $x$  is an independent variable. let  $u_0(x, t)$  denote an initial guess of the exact solution  $u(x, t)$ ,  $h \neq 0$  an auxiliary parameter,  $H_1(x, t) \neq 0$  an auxiliary function, and  $L$  an auxiliary linear operator with the property  $L[r(x, t)] = 0$  when  $r(x, t) = 0$ . Then using  $q \in [0, 1]$  as an embedding parameter, we construct a homotopy as follows:

$$(1-q)L[\phi(x, t; q) - u_0(x, t)] - qhH_1(x, t)N[\phi(x, t; q)] = \hat{H}[\phi(x, t; q); u_0(x, t), H_1(x, t), h, q]. \tag{22}$$

It should be emphasized that we have great freedom to choose the initial guess  $u_0(x, t)$ , the auxiliary linear operator  $L$ , the non-zero auxiliary parameter  $h$ , and the auxiliary function  $H_1(x, t)$ .

Enforcing the homotopy (22) to be zero, i.e.,

$$\hat{H}[\phi(x, t; q); u_0(x, t), H_1(x, t), h, q] = 0, \tag{23}$$

we have the so-called zero-order deformation equation

$$(1-q)L[\phi(x, t; q) - u_0(x, t)] = qhH_1(x, t)N[\phi(x, t; q)]. \tag{24}$$

When  $q = 0$ , the zero-order deformation Eq.(24) becomes

$$\phi(x; 0) = u_0(x, t), \tag{25}$$

and when  $q = 1$ , since  $h \neq 0$  and  $H_1(x, t) \neq 0$ , the zero-order deformation Eq.(24) is equivalent to

$$\phi(x, t; 1) = u(x, t). \tag{26}$$

Thus, according to (25) and (26), as the embedding parameter  $q$  increases from 0 to 1,  $\phi(x, t; q)$  varies continuously from the initial approximation  $u_0(x, t)$  to the exact solution  $u(x, t)$ . Such a kind of continuous variation is called deformation in homotopy [26-29].

Due to Taylor's theorem,  $\phi(x, t; q)$  can be expanded in a power series of  $q$  as follows

$$\phi(x, t; q) = u_0(x, t) + \sum_{m=1}^{\infty} u_m(x, t) q^m, \quad (27)$$

where,

$$u_m(x, t) = \frac{1}{m!} \frac{\partial^m \phi(x, t; q)}{\partial q^m} \Big|_{q=0}.$$

Let the initial guess  $u_0(x, t)$ , the auxiliary linear parameter  $L$ , the nonzero auxiliary parameter  $h$  and the auxiliary function  $H_1(x, t)$  be properly chosen so that the power series (27) of  $\phi(x, t; q)$  converges at  $q = 1$ , then, we have under these assumptions the solution series

$$u(x, t) = \phi(x, t; 1) = u_0(x, t) + \sum_{m=1}^{\infty} u_m(x, t). \quad (28)$$

From Eq.(27), we can write Eq.(24) as follows

$$\begin{aligned} (1-q)L[\phi(x, t, q) - u_0(x, t)] &= (1-q)L\left[\sum_{m=1}^{\infty} u_m(x, t) q^m\right] = q h H_1(x, t) N[\phi(x, t, q)] \Rightarrow \\ L\left[\sum_{m=1}^{\infty} u_m(x, t) q^m\right] - q L\left[\sum_{m=1}^{\infty} u_m(x, t) q^m\right] &= q h H_1(x, t) N[\phi(x, t, q)] \end{aligned} \quad (29)$$

By differentiating (29)  $m$  times with respect to  $q$ , we obtain

$$\begin{aligned} \{L[\sum_{m=1}^{\infty} u_m(x, t) q^m] - q L[\sum_{m=1}^{\infty} u_m(x, t) q^m]\}^{(m)} &= \{q h H_1(x, t) N[\phi(x, t, q)]\}^{(m)} = \\ m! L[u_m(x, t) - u_{m-1}(x, t)] &= h H_1(x, t) m \frac{\partial^{m-1} N[\phi(x, t; q)]}{\partial q^{m-1}} \Big|_{q=0}. \end{aligned}$$

Therefore,

$$L[u_m(x, t) - \chi_m u_{m-1}(x, t)] = h H_1(x, t) \mathfrak{R}_m(u_{m-1}(x, t)), \quad (30)$$

where,

$$\mathfrak{R}_m(u_{m-1}(x, t)) = \frac{1}{(m-1)!} \frac{\partial^{m-1} N[\phi(x, t; q)]}{\partial q^{m-1}} \Big|_{q=0}, \quad (31)$$

and

$$\chi_m = \begin{cases} 0, & m \leq 1 \\ 1, & m > 1 \end{cases}$$

Note that the high-order deformation Eq.(30) is governing the linear operator  $L$ , and the term  $\mathfrak{R}_m(u_{m-1}(x, t))$  can be expressed simply by (31) for any nonlinear operator  $N$ .

To obtain the approximation solution of Eq.(1), according to HAM, let

$$N[u(x, t)] = u(x, t) - G(x, t) + \int_0^t F_1(u(x, t)) dt + \int_0^t F_2(u(x, t)) dt - \int_0^t F_3(u(x, t)) dt,$$

so,

$$\mathfrak{R}_m(u_{m-1}(x, t)) = u_{m-1}(x, t) - G(x, t) + \int_0^t F_1(u_{m-1}(x, t)) dt + \int_0^t F_2(u_{m-1}(x, t)) dt - \int_0^t F_3(u_{m-1}(x, t)) dt. \quad (32)$$

Substituting (32) into (30)

$$L[u_m(x, t) - \chi_m u_{m-1}(x, t)] = hH_1(x, t)[u_{m-1}(x, t) + \int_0^t F_1(u_{m-1}(x, t)) dt + \int_0^t F_2(u_{m-1}(x, t)) dt - \int_0^t F_3(u_{m-1}(x, t)) dt + (1 - \chi_m)G(x, t)]. \quad (33)$$

We take an initial guess  $u_0(x, t) = G(x, t)$ , an auxiliary linear operator  $Lu = u$ , a nonzero auxiliary parameter  $h = -1$ , and auxiliary function  $H_1(x, t) = 1$ . This is substituted into (33) to give the recurrence relation

$$u_0(x, t) = G(x, t), \quad (34)$$

$$u_{n+1}(x, t) = -\int_0^t F_1(u_n(x, t)) dt - \int_0^t F_2(u_n(x, t)) dt + \int_0^t F_3(u_n(x, t)) dt, \quad n \geq 0.$$

Therefore, the solution  $u(x, t)$  becomes

$$u(x, t) = \sum_{n=0}^{\infty} u_n(x, t) \quad (35)$$

$$= G(x, t) + \sum_{n=1}^{\infty} \left( -\int_0^t F_1(u_n(x, t)) dt - \int_0^t F_2(u_n(x, t)) dt + \int_0^t F_3(u_n(x, t)) dt \right).$$

Which is the method of successive approximations. If

$$|u_n(x, t)| < 1,$$

then the series solution (35) convergence uniformly.

## 2.4 Description of the HPM and MHPM

To explain HPM [30-36], we consider the following general nonlinear differential equation:

$$Lu + Nu = f(u), \quad (36)$$

with initial conditions

$$u(x, 0) = f(x).$$

According to HPM, we construct a homotopy which satisfies the following relation

$$H(u, p) = Lu - Lv_0 + p Lv_0 + p [Nu - f(u)] = 0, \quad (37)$$



where  $p \in [0,1]$  is an embedding parameter and  $v_0$  is an arbitrary initial approximation satisfying the given initial conditions.

In HPM, the solution of Eq.(37) is expressed as

$$u(x,t) = u_0(x,t) + p u_1(x,t) + p^2 u_2(x,t) + \dots \quad (38)$$

Hence the approximate solution of Eq.(36) can be expressed as a series of the power of  $p$ , i.e.

$$u = \lim_{p \rightarrow 1} u = u_0 + u_1 + u_2 + \dots$$

where,

$$\begin{aligned} u_0(x,t) &= G(x,t), \\ \vdots \\ u_m(x,t) &= \sum_{k=0}^{m-1} - \int_0^t F_1(u_{m-k-1}(x,t)) dt - \int_0^t F_2(u_{m-k-1}(x,t)) dt + \\ &\quad \int_0^t F_3(u_{m-k-1}(x,t)) dt, \quad m \geq 1. \end{aligned} \quad (39)$$

To explain MHPM [37-42], we consider Eq.(1) as

$$\begin{aligned} L(u) &= u(x,t) - G(x,t) + \int_0^t F_1(u(x,t)) dt + \int_0^t F_2(u(x,t)) dt \\ &\quad - \int_0^t F_3(u(x,t)) dt. \end{aligned}$$

Where  $F_1(u(x,t)) = g_1(x)h_1(t)$ ,  $F_2(u(x,t)) = g_2(x)h_2(t)$  and  $F_3(u(x,t)) = g_3(x)h_3(t)$ . We can define homotopy  $H(u, p, m)$  by

$$H(u, 0, m) = f(u), \quad H(u, 1, m) = L(u),$$

where,  $m$  is an unknown real number and

$$f(u(x,t)) = u(x,t) - z(x,t).$$

Typically we may choose a convex homotopy by

$$H(u, p, m) = (1-p)f(u) + p L(u) + p(1-p)[m(g_1(x) + g_2(x) + g_3(x))] = 0, \quad 0 \leq p \leq 1. \quad (40)$$

Where  $m$  is called the accelerating parameters, and for  $m=0$  we define  $H(u, p, 0) = H(u, p)$ , which is the standard HPM.

The convex homotopy (40) continuously trace an implicitly defined curve from a starting point  $H(u(x,t) - f(u), 0, m)$  to a solution function  $H(u(x,t), 1, m)$ . The embedding parameter  $p$  monotonically increase from 0 to 1 as trivial problem  $f(u) = 0$  is continuously deformed to original problem  $L(u) = 0$ .

The MHPM uses the homotopy parameter  $p$  as an expanding parameter to obtain

$$v = \sum_{n=0}^{\infty} p^n u_n, \quad (41)$$

when  $p \rightarrow 1$ , Eq.(37) corresponds to the original one and Eq.(41) becomes the approximate solution of Eq.(1), i.e.,

$$u = \lim_{p \rightarrow 1} v = \sum_{m=0}^{\infty} u_m.$$

Where,

$$\begin{aligned} u_0(x, t) &= G(x, t), \\ u_1(x, t) &= -\int_0^t F_1(u_0(x, t)) dt - \int_0^t F_2(u_0(x, t)) dt + \int_0^t F_3(u_0(x, t)) dt - m(g_1(x) + g_2(x) + g_3(x)), \\ u_2(x, t) &= -\int_0^t F_1(u_1(x, t)) dt - \int_0^t F_2(u_1(x, t)) dt + \int_0^t F_3(u_1(x, t)) dt + m(g_1(x) + g_2(x) + g_3(x)), \\ &\vdots \\ u_m(x, t) &= \sum_{k=0}^{m-1} -\int_0^t F_1(u_{m-k-1}(x, t)) dt - \int_0^t F_2(u_{m-k-1}(x, t)) dt + \int_0^t F_3(u_{m-k-1}(x, t)) dt, m \geq 3. \end{aligned} \quad (42)$$

### 3 Existence and convergency of iterative methods

We set,

$$\begin{aligned} \alpha_1 &:= T(L_1 + L_2 + L_3), \\ \beta_1 &:= 1 - T(1 - \alpha_1), \quad \gamma_1 := 1 - T\alpha_1. \end{aligned}$$

**Theorem 3.1** Let  $0 < \alpha_1 < 1$ , then Kadomtsev-Petviashvili equation (1), has a unique solution.

**Proof.** Let  $u$  and  $u^*$  be two different solutions of (3) then

$$\begin{aligned} |u - u^*| &= \left| -\int_0^t [F_1(u(x, t)) - F_1(u^*(x, t))] dt - \int_0^t [F_2(u(x, t)) - F_2(u^*(x, t))] dt \right. \\ &\quad \left. + \int_0^t [F_3(u(x, t)) - F_3(u^*(x, t))] dt \right| \leq \int_0^t |F_1(u(x, t)) - F_1(u^*(x, t))| dt + \int_0^t |F_2(u(x, t)) - F_2(u^*(x, t))| dt + \\ &\quad \int_0^t |F_3(u(x, t)) - F_3(u^*(x, t))| dt \leq T(L_1 + L_2 + L_3) |u - u^*| = \alpha_1 |u - u^*|. \end{aligned}$$

From which we get  $(1 - \alpha_1) |u - u^*| \leq 0$ . Since  $0 < \alpha_1 < 1$ , then  $|u - u^*| = 0$ . Implies  $u = u^*$  and completes the proof.

**Theorem 3.2** The series solution  $u(x, t) = \sum_{i=0}^{\infty} u_i(x, t)$  of problem(1) using MADM convergence when

$$0 < \alpha_1 < 1, \quad |u_1(x, t)| < \infty.$$

**Proof.** Denote as  $(C[J], \|\cdot\|)$  the Banach space of all continuous functions on  $J$  with the norm  $\|G(x, t)\| = \max |G(x, t)|$ , for all  $x, t$  in  $J$ . Define the sequence of partial sums  $s_n$ , let  $s_n$  and  $s_m$  be arbitrary partial sums with  $n \geq m$ . We are going to prove that  $s_n$  is a Cauchy sequence in this Banach space:

$$\begin{aligned} \|s_n - s_m\| &= \max_{\forall t \in J} |s_n - s_m| = \max_{\forall t \in J} \left| \sum_{i=m+1}^n u_i(x, t) \right| = \\ &= \max_{\forall t \in J} \left| - \int_0^t \left( \sum_{i=m}^{n-1} A_i \right) dt - \int_0^t \left( \sum_{i=m}^{n-1} B_i \right) dt + \int_0^t \left( \sum_{i=m}^{n-1} L_i \right) dt \right|. \end{aligned}$$

From [15], we have

$$\begin{aligned} \sum_{i=m}^{n-1} A_i &= F_1(s_{n-1}) - F_1(s_{m-1}), \\ \sum_{i=m}^{n-1} B_i &= F_2(s_{n-1}) - F_2(s_{m-1}), \\ \sum_{i=m}^{n-1} L_i &= F_3(s_{n-1}) - F_3(s_{m-1}). \end{aligned}$$

So,

$$\begin{aligned} \|s_n - s_m\| &= \max_{\forall t \in J} \left| - \int_0^t [F_1(s_{n-1}) - F_1(s_{m-1})] dt - \int_0^t [F_2(s_{n-1}) - F_2(s_{m-1})] dt + \int_0^t [F_3(s_{n-1}) - F_3(s_{m-1})] dt \right| \leq \\ &= \int_0^t |F_1(s_{n-1}) - F_1(s_{m-1})| dt + \int_0^t |F_2(s_{n-1}) - F_2(s_{m-1})| dt \\ &+ \int_0^t |F_3(s_{n-1}) - F_3(s_{m-1})| dt \leq \alpha_1 \|s_n - s_m\|. \end{aligned}$$

Let  $n = m + 1$ , then

$$\|s_n - s_m\| \leq \alpha_1 \|s_m - s_{m-1}\| \leq \alpha_1^2 \|s_{m-1} - s_{m-2}\| \leq \dots \leq \alpha_1^m \|s_1 - s_0\|.$$

From the triangle inequality we have

$$\begin{aligned} \|s_n - s_m\| &\leq \|s_{m+1} - s_m\| + \|s_{m+2} - s_{m+1}\| + \dots + \|s_n - s_{n-1}\| \leq [\alpha_1^m + \alpha_1^{m+1} + \dots + \alpha_1^{n-m-1}] \|s_1 - s_0\| \\ &\leq \alpha_1^m [1 + \alpha_1 + \alpha_1^2 + \dots + \alpha_1^{n-m-1}] \|s_1 - s_0\| \leq \alpha_1^m \left[ \frac{1 - \alpha_1^{n-m}}{1 - \alpha_1} \right] \|u_1(x, t)\|. \end{aligned}$$

Since  $0 < \alpha_1 < 1$ , we have  $(1 - \alpha_1^{n-m}) < 1$ , then

$$\|s_n - s_m\| \leq \frac{\alpha_1^m}{1 - \alpha_1} \max_{\forall t \in J} |u_1(x, t)|. \quad (43)$$

But  $|u_1(x, t)| < \infty$ , so, as  $m \rightarrow \infty$ , then  $\|s_n - s_m\| \rightarrow 0$ . We conclude that  $s_n$  is a Cauchy sequence in  $C[J]$ , therefore the series is convergence and the proof is complete.

**Theorem 3.3** The maximum absolute truncation error of the series solution

$u(x, t) = \sum_{i=0}^{\infty} u_i(x, t)$  to problem (1) by using MADM is estimated to be

$$\max |u(x, t) - \sum_{i=0}^m u_i(x, t)| \leq \frac{k\alpha_1^m}{1-\alpha_1}. \quad (44)$$

**Proof.** From inequality (43), when  $n \rightarrow \infty$ , then  $s_n \rightarrow u$  and

$$\max |u_1(x, t)| \leq T(\max_{\forall t \in J} |F_1(u_0(x, t))| + \max_{\forall t \in J} |F_2(u_0(x, t))| + \max_{\forall t \in J} |F_3(u_0(x, t))|).$$

Therefore,

$$\|u(x, t) - s_m\| \leq \frac{\alpha_1^m}{1-\alpha_1} T(\max_{\forall t \in J} |F_1(u_0(x, t))| + \max_{\forall t \in J} |F_2(u_0(x, t))| + \max_{\forall t \in J} |F_3(u_0(x, t))|).$$

Finally the maximum absolute truncation error in the interval  $J$  is obtained by (44).

**Theorem 3.4** The solution  $u_n(x, t)$  obtained from the relation (20) using VIM converges to the exact solution of the problem (1) when  $0 < \alpha_1 < 1$  and  $0 < \beta_1 < 1$ .

**Proof.**

$$u_{n+1}(x, t) = u_n(x, t) - L_t^{-1}([u_n(x, t) - G(x, t) + \int_0^t F_1(u_n(x, t)) dt + \int_0^t F_2(u_n(x, t)) dt - \int_0^t F_3(u_n(x, t)) dt]), \quad (45)$$

$$u(x, t) = u(x, t) - L_t^{-1}([u(x, t) - G(x, t) + \int_0^t F_1(u(x, t)) dt + \int_0^t F_2(u(x, t)) dt - \int_0^t F_3(u(x, t)) dt]). \quad (46)$$

By subtracting relation (45) from (46),

$$u_{n+1}(x, t) - u(x, t) = u_n(x, t) - u(x, t) - L_t^{-1}(u_n(x, t) - u(x, t) + \int_0^t [F_1(u_n(x, t)) - F_1(u(x, t))] dt + \int_0^t [F_2(u_n(x, t)) - F_2(u(x, t))] dt - \int_0^t [F_3(u_n(x, t)) - F_3(u(x, t))] dt),$$

if we set,  $e_{n+1}(x, t) = u_{n+1}(x, t) - u_n(x, t)$ ,  $e_n(x, t) = u_n(x, t) - u(x, t)$ ,  $|e_n(x, t^*)| = \max_t |e_n(x, t)|$  then since  $e_n$  is a decreasing function with respect to  $t$  from the mean value theorem we can write,

$$\begin{aligned}
e_{n+1}(x, t) &= e_n(x, t) + L_t^{-1}(-e_n(x, t) - \int_0^t [F_1(u_n(x, t)) - F_1(u(x, t))] dt \\
&\quad - \int_0^t [F_2(u_n(x, t)) - F_2(u(x, t))] dt + r \int_0^t [F_3(u_n(x, t)) - F_3(u(x, t))] dt) \\
&\leq e_n(x, t) + L_t^{-1}[-e_n(x, t) + L_t^{-1} |e_n(x, t)| (T(L_1 + L_2 + L_3))] \\
&\leq e_n(x, t) - T e_n(x, \eta) + T(L_1 + L_2 + L_3) L_t^{-1} L_t^{-1} |e_n(x, t)| \\
&\leq (1 - T(1 - \alpha_1)) |e_n(x, t^*)|,
\end{aligned}$$

where  $0 \leq \eta \leq t$ . Hence,  $e_{n+1}(x, t) \leq \beta_1 |e_n(x, t^*)|$ .

Therefore,

$$\|e_{n+1}\| = \max_{t \in J} |e_{n+1}| \leq \beta_1 \max_{t \in J} |e_n| \leq \beta_1 \|e_n\|.$$

Since  $0 < \beta_1 < 1$ , then  $\|e_n\| \rightarrow 0$ . So, the series converges and the proof is complete.

**Theorem 3.5** The solution  $u_n(x, t)$  obtained from the relation (22) using MVIM for the problem (1) converges when  $0 < \alpha_1 < 1$ ,  $0 < \gamma_1 < 1$ .

**Proof.** The Proof is similar to the previous theorem.

**Theorem 3.6** The maximum absolute truncation error of the series solution

$u(x, t) = \sum_{i=0}^{\infty} u_i(x, t)$  to problem (1) by using VIM is estimated to be

$$\|e_n\| \leq \frac{\beta_1^n k'}{1 - \beta_1}, \quad k' = \max |u_1(x, t)|.$$

**Proof.**

$$\begin{aligned}
u_{n+1} - u_n &= (u_{n+1} - u) + (u - u_n) = e_n - e_{n+1} \\
\rightarrow e_n &= e_{n+1} - (u_{n+1} - u_n) \\
\|e_n\| &= \|e_{n+1} - (u_{n+1} - u_n)\| \leq \|e_{n+1}\| + \|u_{n+1} - u_n\| \leq \beta_1 \|e_n\| + \|u_{n+1} - u_n\| \\
\rightarrow \|e_n\| &\leq \frac{\|u_{n+1} - u_n\|}{1 - \beta_1} \leq \frac{\beta_1^n k'}{1 - \beta_1}.
\end{aligned}$$

**Theorem 3.7** If the series solution (34) of problem (1) using HAM convergent then it converges to the exact solution of the problem (1).

**Proof.** We assume:

$$\begin{aligned}
u(x, t) &= \sum_{m=0}^{\infty} u_m(x, t), \\
\hat{F}_1(u(x, t)) &= \sum_{m=0}^{\infty} F_1(u_m(x, t)), \\
\hat{F}_2(u(x, t)) &= \sum_{m=0}^{\infty} F_2(u_m(x, t)), \\
\hat{F}_3(u(x, t)) &= \sum_{m=0}^{\infty} F_3(u_m(x, t)).
\end{aligned}$$

Where,

$$\lim_{m \rightarrow \infty} u_m(x, t) = 0.$$

We can write,

$$\sum_{m=1}^n [u_m(x, t) - \chi_m u_{m-1}(x, t)] = u_1 + (u_2 - u_1) + \dots + (u_n - u_{n-1}) = u_n(x, t). \quad (47)$$

Hence, from (47),

$$\lim_{n \rightarrow \infty} u_n(x, t) = 0. \quad (48)$$

So, using (48) and the definition of the linear operator  $L$ , we have

$$\sum_{m=1}^{\infty} L[u_m(x, t) - \chi_m u_{m-1}(x, t)] = L\left[\sum_{m=1}^{\infty} [u_m(x, t) - \chi_m u_{m-1}(x, t)]\right] = 0.$$

therefore from (30), we can obtain that,

$$\sum_{m=1}^{\infty} L[u_m(x, t) - \chi_m u_{m-1}(x, t)] = hH_1(x, t) \sum_{m=1}^{\infty} \mathfrak{R}_{m-1}(u_{m-1}(x, t)) = 0.$$

Since  $h \neq 0$  and  $H_1(x, t) \neq 0$ , we have

$$\sum_{m=1}^{\infty} \mathfrak{R}_{m-1}(u_{m-1}(x, t)) = 0. \quad (49)$$

By substituting  $\mathfrak{R}_{m-1}(u_{m-1}(x, t))$  into the relation (49) and simplifying it, we have

$$\begin{aligned} \sum_{m=1}^{\infty} \mathfrak{R}_{m-1}(u_{m-1}(x, t)) &= \sum_{m=1}^{\infty} \left[ \int_0^t F_1(u_{m-1}(x, t)) dt \right. \\ &+ \int_0^t F_2(u_{m-1}(x, t)) dt - \int_0^t F_3(u_{m-1}(x, t)) dt + (1 - \chi_m)G(x, t)(x) \bigg] \\ &= u(x, t) - G(x, t) + \int_0^t \hat{F}_1(u(x, t)) dt + \int_0^t \hat{F}_2(u(x, t)) dt - \int_0^t \hat{F}_3(u(x, t)) dt. \end{aligned} \quad (50)$$

From (49) and (50), we have

$$u(x, t) = G(x, t) - \int_0^t \hat{F}_1(u(x, t)) dt - \int_0^t \hat{F}_2(u(x, t)) dt + \int_0^t \hat{F}_3(u(x, t)) dt.$$

Therefore,  $u(x, t)$  must be the exact solution.

**Theorem 3.8** The maximum absolute truncation error of the series solution

$u(x, t) = \sum_{i=0}^{\infty} u_i(x, t)$  to problem (1) by using HAM is estimated to be

$$\|e_n\| \leq \frac{\alpha_1^n k'}{1 - \alpha_1}, \quad k' = \max |u_1(x, t)|.$$

**Proof.** The Proof is similar to the 3.6 theorem

**Theorem 3.9** If  $|u_m(x, t)| \leq 1$ , then the series solution  $u(x, t) = \sum_{i=0}^{\infty} u_i(x, t)$  of problem (1) converges to the exact solution by using HPM.

**Proof.** We set,

$$\begin{aligned} \phi_n(x, t) &= \sum_{i=1}^n u_i(x, t), \\ \phi_{n+1}(x, t) &= \sum_{i=1}^{n+1} u_i(x, t). \\ |\phi_{n+1}(x, t) - \phi_n(x, t)| &= D(\phi_{n+1}(x, t), \phi_n(x, t)) = D(\phi_n + u_n, \phi_n) \\ &= D(u_n, 0) \leq \sum_{k=0}^{m-1} \int_0^t |F_1(u_{m-k-1}(x, t))| dt + \int_0^t |F_2(u_{m-k-1}(x, t))| dt \\ &\quad + \int_0^t |F_3(u_{m-k-1}(x, t))| dt. \\ \rightarrow \sum_{n=0}^{\infty} |\phi_{n+1}(x, t) - \phi_n(x, t)| &\leq m\alpha_1 |G(x, t)| \sum_{n=0}^{\infty} (m\alpha_1)^n. \end{aligned}$$

Therefore,

$$\lim_{n \rightarrow \infty} u_n(x, t) = u(x, t).$$

**Theorem 3.10** If  $|u_m(x, t)| \leq 1$ , then the series solution  $u(x, t) = \sum_{i=0}^{\infty} u_i(x, t)$  of problem (1) converges to the exact solution by using MHPM.

**Proof.** The Proof is similar to the previous theorem.

**Theorem 3.11** The maximum absolute truncation error of the series solution

$u(x, t) = \sum_{i=0}^{\infty} u_i(x, t)$  to problem (1) by using HPM is estimated to be

$$\|e_n\| \leq \frac{(n\alpha_1)^n nk'}{1 - \alpha_1}, \quad k' = \max |u_1(x, t)|.$$

**Proof.** The Proof is similar to the 3.6 theorem

## 4 Numerical example

In this section, we compute a numerical example which is solved by the ADM, MADM, VIM,

MVIM, HPM, MHPM and HAM. The program has been provided with Mathematica 6 according to the following algorithm where  $\varepsilon$  is a given positive value.

**Algorithm 1:**

**Step 1.** Set  $n \leftarrow 0$ .

**Step 2.** Calculate the recursive relations (10) for ADM, (13) for MADM, (34) for HAM, (39) for HPM and (42) for MHPM.

**Step 3.** If  $|u_{n+1} - u_n| < \varepsilon$  then go to step 4,  
else  $n \leftarrow n + 1$  and go to step 2.

**Step 4.** Print  $u(x, t) = \sum_{i=0}^n u_i(x, t)$  as the approximate of the exact solution.

**Algorithm 2:**

**Step 1.** Set  $n \leftarrow 0$ .

**Step 2.** Calculate the recursive relations (20) for VIM and (21) for MVIM.

**Step 3.** If  $|u_{n+1} - u_n| < \varepsilon$  then go to step 4,  
else  $n \leftarrow n + 1$  and go to step 2.

**Step 4.** Print  $u_n(x, t)$  as the approximate of the exact solution.

**Example 4.1** Consider the Kadomtsev-Petviashvili equation as follows:

$$u_t(x, t) + (x + t)u_x(x, t) + \frac{1}{2}u_{xx}(x, t) - u(x, t) = 0.$$

With initial condition:

$$g(x) = e^x.$$

**Table 1** Numerical results for Example 4.1

(x,t)	Errors			
	ADM(n=16)	MADM(n=13)	VIM(n=9)	MVIM(n=8)
(0.1,0.15)	0.081436	0.073123	0.050659	0.043416
(0.2,0.17)	0.082589	0.074356	0.051375	0.044237
(0.3,0.20)	0.083296	0.075419	0.051842	0.044732
(0.4,0.23)	0.083746	0.075729	0.052321	0.045144
(0.5,0.25)	0.084315	0.076348	0.052796	0.045748
(0.7,0.30)	0.085228	0.076808	0.053225	0.046207
(0.9,0.35)	0.086708	0.077173	0.053705	0.046875
(1.0,0.40)	0.087417	0.077783	0.054202	0.047089



(x,t)	Errors		
	HPM(n=9)	MHPM(n=8)	HAM(n=5)
(0.1,0.15)	0.062156	0.033646	0.022544
(0.20,0.17)	0.063015	0.034349	0.023154
(0.3,0.20)	0.063385	0.035172	0.023557
(0.4,0.23)	0.065317	0.035471	0.024048
(0.5,0.25)	0.065819	0.036163	0.024829
(0.7,0.30)	0.066112	0.036372	0.025362
(0.9,0.35)	0.066443	0.36724	0.026315
(1.0,0.40)	0.067236	0.037052	0.026759

Table 1, shows that, approximate solution of the Kadomtsev-Petviashvili equation is convergence with 5 iterations by using the HAM . By comparing the results of Table 1 , we can observe that the HAM is more rapid convergence than the ADM, MADM, VIM, MVIM, HPM and MHPM.

## 5 Conclusions

The homotopy analysis method has been known as a powerful scheme for solving many functional equations such as algebraic equations, ordinary and partial differential equations, integral equations and so on. The HAM has been shown to solve effectively, easily and accurately a large class of nonlinear problems with approximations converge rapidly to accurate the solution. In this work, the HAM has been successfully employed to obtain the approximate solution of the Kadomtsev-Petviashvili equation. We showed that the homotopy analysis method has more rapid convergence than the ADM, MADM, MVIM, HPM , MHPM and VIM.

## References

1. Kadomtsev,B., Petviashvili,P.,(1970). On the stability of solitary waves in weakly dispersive media.Sov. Phys. Dokl,15,539–541.
2. Hu,Y., Ma,J.,(2004). Nonlinear Kadomtsev-Petviashvili equation and discrete- functional- type BSDEs with continuous coefficients.Stochastic Processes and their Applications, 112, 23-51.
3. Morette,C.D., Zhang,.T.R.,(1983). Kadomtsev-Petviashvili equation in phase with application to coherent-state transitions. Physical Review D, 28 ,2517-2525.
4. Takeda, M., (1998). Asymptotic properties of generalized Kadomtsev-Petviashvili equation.Potential Analysis, 9 ,261-291.
5. Kwas, M., Li,Y.,(2003). Worst case complexity of multivariate Kadomtsev-Petviashvili equation integration. Journal of Complexity, 19,730-743.
6. Bond,.S.D., Laird, B.B., Leimkuhler,B.J.,(2003). On the approximation of Kadomtsev-Petviashvili path integrals. Journal of Computational Physics, 185,472-483.
7. Buchmann,F.M., Petersen,W.D.,(2003). Solving dirichlet problems numerically using the Kadomtsev-Petviashvili equation .BIT Numerical Mathematics, 43, 519-540.
8. Ouerdiane,H., Silva,J.L.,(202). Generalized Kadomtsev-Petviashvili equation with stochastic potential. In finite Dimensional Analysis.Quantum Probability and Related Topics, 5, 243-255.
9. Ostrovsky, D.,(2007). Functional Kadomtsev-Petviashvili equation for limit lognormal multifractals. Journal of statistical physics, 127 , 935-965.

10. Turgeman, L., Carmi, S., Barkai, E., (2009). Fractional Kadomtsev-Petviashvili equation for Non-Brownian functionals. *Physical Review Letters*, 103, 1-4.
11. Behriy, S.H., Hashish, H., E-Kalla, I.L., Elsaid, A., (2007). A new algorithm for the decomposition solution of nonlinear differential equations. *App.Math.Comput*, 54, 459-466.
12. Fariborzi Araghi, M.A., Behzadi, Sh.S., (2009). Solving nonlinear Volterra-Fredholm integral differential equations using the modified Adomian decomposition method. *Comput. Methods in Appl. Math*, 9, 1-11.
13. Wazwaz, A.M., (2001). Construction of solitary wave solution and rational solutions for the KdV equation by ADM. *Chaos, Solution and fractals*, 12, 2283-2293.
14. Wazwaz, A.M., (1997). A first course in integral equations, WSPC, New Jersey.
15. El-Kalla, I.L., (2008). Convergence of the Adomian method applied to a class of nonlinear integral equations, *Appl.Math.Comput*, 21, 372-376.
16. He, J.H., Wu, X.H., (2006). Exp-function method for nonlinear wave equations. *Chaos, Solitons and Fractals*, 30, 700-708.
17. He, J.H., (2004). Variational principle for some nonlinear partial differential equations with variable coefficients. *Chaos, Solitons and Fractals*, 19, 847-851.
18. He, J.H., Shu-Qiang, W., (2007). Variational iteration method for solving integro-differential equations. *Physics Letters A*, 367, 188-191.
19. He, J.H., (2007). Variational iteration method some recent results and new interpretations. *J. Comp. and Appl. Math*, 207, 3-17.
20. Fariborzi Araghi, M.A., Behzadi, Sh.S., (2010). Solving nonlinear Volterra-Fredholm integro-differential equations using He's variational iteration method. *International Journal of Computer Mathematics*, DOI: 10.1007/s12190-010-0417-4.
21. Yildirim, A., (2010). Variational iteration method for modified Camassa-Holm and Degasperis-Procesi equations. *International Journal for Numerical Methods in Biomedical Engineering*, 26, 266-272.
22. Abbasbandy, S., (2008). Numerical method for non-linear wave and diffusion equations by the variational iteration method. *Int. J. Numer. Methods Eng.* 73, 1836-1843.
23. Behzadi, Sh.S., Fariborzi Araghi, M.A., (2011). The use of iterative methods for solving Navier-Stokes equation. *J.Appl.Math.Informatics*, 29, 1-15.
24. Abassy, T.A., Tawil, E.L., Zoheiry, H.E.L., (2007). Toward a modified variational iteration method (MVIM). *J.Comput.Appl.Math*, 207, 137-147.
25. Abassy, T.A., Tawil, E.L., Zoheiry, H.E.L., (2007). Modified variational iteration method for Boussinesq equation. *Comput.Math.Appl*, 54, 955-965.
26. Liao, S.J., (2003). *Beyond Perturbation: Introduction to the Homotopy Analysis Method*. Chapman and Hall/CRC Press, Boca Raton.
27. Liao, S.J., (2009). Notes on the homotopy analysis method: some definitions and theorems. *Communication in Nonlinear Science and Numerical Simulation*, 14, 983-997.
28. Fariborzi Araghi, M.A., Behzadi, Sh.S., (2010). Numerical solution of nonlinear Volterra-Fredholm integro-differential equations using Homotopy analysis method. *Journal of Applied Mathematics and Computing*, DOI: 10.1080/00207161003770394.
29. Babolian, E., Saeidian, J., (2009). Analytic approximate solutions to Burger, Fisher, Huxley equations and two combined forms of these equations. *Commun Nonlinear Sci Numer Simulat*, 14, 1984-1992.
30. Biazar, J., Ghazvini, H., (2009). Convergence of the homotopy perturbation method for partial differential equations. *Nonlinear Analysis: Real World Application*, 10, 2633-2640.
31. Ghasemi, M., Tavasoli, M., Babolian, E., (2007). Application of He's homotopy perturbation method of nonlinear integro-differential equation. *Appl.Math.Comput*, 188, 538-548.
32. Golbabai, A., Keramati, B., (2009). Solution of non-linear Fredholm integral equations of the first kind using modified homotopy perturbation method. *Chaos Solitons and Fractals*, 5, 2316-2321.
33. Javidi, M., (2009). Modified homotopy perturbation method for solving linear Fredholm integral equations. *Chaos Solitons and Fractals*, 50, 159-165.
34. Fariborzi Araghi, M.A., Behzadi, Sh.S., (2011). Numerical solution for solving Burger's-Fisher equation by using Iterative Methods. *Mathematical and Computational Applications*, 16, 443-455.
35. Abbasbandy, S., (2006). Modified homotopy perturbation method for nonlinear equations and comparison with Adomian decomposition method. *Appl.Math.Comput* 172, 431-438.
36. Yildirim, A., Mohyud-Din, S.T., Zhang, D.H., (2010). Analytical solutions to the pulsed Klein-Gordon equation using Modified Variational Iteration Method (MVIM) and Boubaker Polynomials Expansion Scheme (BPES). *Computers and Mathematics with Applications*, 59, 2473-2477.
37. Sezer, S.A., Yildirim, A., Mohyud-Din, S.T., (2011). He's homotopy perturbation method for solving the fractional KdV-Burgers-Kuramoto equation. *International Journal of Numerical Methods for Heat and Fluid*

- Flow, 21,448-458.
38. Yildirim,A.,(2008). Solution of BVPs for Fourth-Order Integro-Differential Equations by using Homotopy Perturbation Method. Computers and Mathematics with Applications, 56,3175-3180.
  39. Yildirim,A.,(2008). The Homotopy Perturbation Method for Approximate Solution of the Modified KdV Equation. Zeitschrift für Naturforschung A,A Journal of Physical Sciences, 63,621-626.
  40. Yildirim,A.,(2010). Application of the Homotopy perturbation method for the Fokker-Planck equation.International Journal for Numerical Methods in Biomedical Engineering, 26,1144-1154.
  41. Behzadi,Sh.S.,(2010). The convergence of homotopy methods for nonlinear Klein-Gordon equation. J.Appl.Math.Informatics, 28,1227-1237.
  42. Abbasbandy,S. ,Shirzadi,A., (2008).The variational iteration method for a class of eight-order boundary value differential equations. Z. Naturforsch. 63,745-751.